

**Fort Walton Beach**

1034 Mar Walt Drive  
Fort Walton Beach,  
FL 32547

**Destin**

36474C Emerald  
Coast Parkway, Suite 3101  
Destin, FL 32541

**Niceville**

554-D Twin  
Cities Boulevard  
Niceville, FL 32578

**Panama City**

1827 Harrison Avenue  
Panama City, FL 32405

**Crestview**

5300 South Fardon Boulevard  
Crestview, FL 32536

P 850-863-2153 F 850-809-4312

[www.orthoassociates.net](http://www.orthoassociates.net)

## Dr. Landry ALIF/XLIF/SL Joint Fusion Instructions / Information

**The Surgery Center  
at Sacred Heart Medical Park**  
36500 Emerald Coast Parkway  
Destin, FL 32541  
(850) 269-0301

**Sacred Heart Hospital  
on the Emerald Coast**  
7800 Emerald Coast Pkwy  
Destin, FL 32550  
(850) 278-3000

**Fort Walton Beach Medical Center**  
1000 Mar Walt Drive  
Fort Walton Bch, FL 32547  
(850) 862-1111

**Surgical Specialists, ASC**  
1034 Mar Walton Drive  
Fort Walton Bch, FL 32547  
(850) 315-9385

- 1. NO SMOKING!** There are numerous very well-designed studies that demonstrate nicotine indirectly inhibits and prevents bone healing and bony fusion from occurring. Not only does nicotine use cause blood vessel constriction, it also has a direct effect on the bone healing process itself. This means no cigarettes, no electronic cigarettes, no chewing tobacco, no nicotine patches, and no nicotine gum. It is the nicotine that is the problem. If you are a smoker, you should immediately discontinue nicotine use as this is the most important thing you can do to prevent a poor outcome. Chantix and Wellbutrin are some medications that have been used to assist with smoking cessation, however, these need to be initiated on a case-by-case basis.
- 2. Blood Thinners.** Blood thinner medications (anti-coagulants) should be stopped 7-10 days before surgery. Examples of anti-coagulants include Plavix, Lovenox, Heparin, Xarelto, Pradaxa, Aspirin, etc. Prior to discontinuing these medications, please speak with your primary care physician, or cardiologist to ensure it is safe to do so. You can restart the prescribed anti-coagulant medications 3 days after the surgery.
- 3. Anti-inflammatories.** You should not take any non-steroidal anti-inflammatory medications (NSAIDs) 10 days prior to surgery and during the first 12 months following surgery. These medications can act as blood thinners and can make the case significantly more difficult to perform if they have been taken within a week of surgery. NSAIDs include Motrin, Ibuprofen, Advil, Aleve, Naproxen, Naprosyn, Celebrex, Vioxx, Etodolac, Lodine, Diclofenac, Meloxicam and Mobic. Well-designed studies have shown that these medications can decrease and even prevent bone healing and bony fusions from occurring. Other medications such as steroids should be avoided, if possible, but should only be discontinued at the discretion of your primary care physician.
- 4. All Other Medications.** All other medications should be discontinued the day of surgery, except for beta blockers. These medications have been found by well-designed studies to be cardioprotective in the acute postoperative time frame. These medications commonly end in – lol, i.e: Metoprolol, labetalol, etc.

5. **Vitamins and Herbal Supplements.** All vitamins and herbal supplements (fish oil, St. John's wort, Multivitamins, etc.) should be discontinued at least 7-10 days prior to surgery and should not be restarted until after the first postoperative visit.
6. **No Bending or Heavy Lifting.** Do not bend or lift anything heavier than 10 pounds for 12 weeks after surgery.
7. **No Twisting.** Do not twist at the waist for 12 weeks after surgery.
8. **The Postoperative Dressing.** The dressing that was applied at the time of the surgery is impregnated with silver and has anti-microbial properties. After 2 weeks, replace with a new dressing and keep it on until the first post-operative appointment. Underneath the dressing will be surgical glue. **Do not remove the surgical glue** – let this naturally fall off.
9. **Showers.** It is okay to take showers beginning on the day after surgery, however, do not take baths or submerge the wound(s). The incision should not be vigorously scrubbed. The incision area can be washed with a clean soapy cloth prior to washing the rest of the body. Pat-dry the surgical wound with a clean, dry towel prior to drying off the rest of the body.
10. **Walking.** You should walk at least 1 mile a day, every day by the 3-week postoperative mark. This can be done in one session, or it can be broken up over multiple sessions over the course of the day. The goal is to reduce the postoperative incidents of deep venous thromboses (DVT) and venous thromboembolism and improve the aerobic capacity of the patient.
11. **Contact Orthopaedic Associates Immediately at (850) 863-2153, if the following occurs:**
  - A persistent fever develops and is greater than 101.5.
  - Uncontrolled pain despite taking pain medications as instructed.
  - Gross purulence (pus) drainage from the wound.
12. **Follow-up with Dr. Landry.** You should follow-up with Dr. Landry approximately 3 weeks after your surgery. The purpose of this visit is to evaluate the wound, evaluate your overall level of pain, and taper the pain medication. Pain medications and muscle relaxers are to be taken only as needed.
13. **Postoperative Visits.** You will be seen, postoperatively, at the following at the following times: 3 weeks, 8 weeks, 3 months, 6 months, and 1 year.
14. **Physical Therapy.** Sometimes, muscular weakness or neurological symptoms take a while to improve. However, if symptoms have not significantly improved by 8 weeks out, or there is significant postoperative stiffness, we will consider initiating physical therapy. If the patient has a job that requires moderate to heavy lifting, we will usually allow the patient to return to work before 8 weeks postoperatively. Alternatively, we will initiate physical therapy at the 8-

**Fort Walton Beach**

1034 Mar Walt Drive  
Fort Walton Beach,  
FL 32547

**Destin**

36474C Emerald  
Coast Parkway, Suite 3101  
Destin, FL 32541

**Niceville**

554-D Twin  
Cities Boulevard  
Niceville, FL 32578

**Panama City**

1827 Harrison Avenue  
Panama City, FL 32405

**Crestview**

5300 South Ferdon Boulevard  
Crestview, FL 32536

P 850-863-2153 F 850-809-4312

[www.orthoassociates.net](http://www.orthoassociates.net)

week postoperative mark and focus on a work hardening program to get the patient ready to go back to work by approximately 3 months postoperatively.

15. **Be Patient.** The primary goal of the surgery is to prevent the patient from getting worse, however, it is common for patients to notice significant improvement, sometimes as early as on the recovery room table. Sometimes it can take upwards of 9-12 months to get significant resolution of the symptoms.
16. **After Surgery,** if you run low on pain medications, you must allow 72 hours for a refill. DO NOT wait until you are out of medication to call us. **The prescription request line is (850) 315-9241.**
17. If you have an illness prior to surgery (within the week of surgery) please notify our office, as your surgery may need to be rescheduled.
18. If an outpatient procedure is scheduled, please arrange to have someone with you at the Surgery Center, drive you home and stay with you for 24 hours following the surgery. You must have a ride to and from the procedure by a responsible adult, friend, or family member. **You CANNOT use taxis, a ridesharing service (Uber, Lyft, etc), or public transportation after surgery. We will not accept the liability, and your case will be cancelled.**
19. For urgent issues after hours or on weekends, please call **(850) 863-2153** and the answering service will contact the physician on-call. **Do not leave a message.** Wait for the prompts, and you will be connected to the answering service.
20. Please bring your insurance information, a list of all prescription medications, and a picture ID with you the day of surgery.
21. If you cancel, or do not show up to one of the appointments coordinated for you before your surgery, it may result in your surgery being cancelled.

DATE OF SURGERY: \_\_\_\_\_

FACILITY: \_\_\_\_\_

ARRIVAL TIME: \_\_\_\_\_

( ) THE FACILITY WILL CALL YOU 24-48 HOURS PRIOR TO SURGHERY WITH YOUR ARRIVAL TIME